

Evidence Summary

Alcohol and Pregnancy

During pregnancy, the baby is vulnerable to exposure to harmful substances including alcohol. Drinking alcohol during pregnancy increases the risk of harm to both mother and baby. These risks include pregnancy complications such as miscarriage, stillbirth, preterm birth, growth restriction (including low birth weight and small for gestational age), and developmental deficits. Fetal Alcohol Spectrum Disorder (FASD) is the term used to describe the range of effects that can occur on the brain and body of babies exposed to alcohol in the womb.

How does alcohol affect the fetus and mother?

Alcohol passes from the mother through the placenta to the fetus at the same concentration as in the mother's blood. But the fetus is unable to process alcohol as efficiently as the mother and the alcohol stays in the fetus' system for longer, having a sustained and prolonged effect¹.

Alcohol interferes with fetal development through direct cell damage or by interfering with cell development². Exposure to alcohol may cause major congenital abnormalities or functional defects to organs. This includes changes in brain development³, cognitive deficits, and behavioural deficits that can have life-long effects⁴⁻⁷.

Fetal Alcohol Spectrum Disorder

*"Fetal Alcohol Spectrum Disorder (FASD) is a diagnostic term used to describe impacts on the brain and body of individuals prenatally exposed to alcohol. FASD is a lifelong disability. Individuals with FASD will experience some degree of challenges in their daily living, and need support with motor skills, physical health, learning, memory, attention, communication, emotional regulation, and social skills to reach their full potential. Each individual with FASD is unique and has areas of both strengths and challenges"*⁸.

There is no data on how common FASD is in New Zealand. However, based on international studies, the Ministry of Health estimates that three in every 100 births may be affected by alcohol - an estimated 1,800 each year⁹.

Alcohol exposure can cause low birth weight^{10 11}, which in turn is associated with poorer long term health outcomes. There is also an increased risk to the mother when drinking during pregnancy including miscarriage, still-birth or preterm birth and their associated complications^{10 12 13}.

What is the level of alcohol use during pregnancy in New Zealand?

One in five women report drinking alcohol during their pregnancy¹⁴, while 13% continue to drink after the first trimester¹⁵.

Drinking more than four drinks per week in the first trimester is more common in younger women (under 19), Māori women, women who smoke, women with fewer education qualifications, and women whose pregnancy is unplanned¹⁶.

However, drinking at lower levels throughout pregnancy is more common in older women (over 40), New Zealand European women, and women from high income households (over \$150,000 per year)¹⁶. These findings highlight the importance of asking all women about their alcohol use during pregnancy.

Are there safe levels and times for consuming alcohol when pregnant?

The Ministry of Health, the Health Promotion Agency /Te Hiringa Hauora, the Royal New Zealand College of General Practitioners, the New Zealand College of Midwives and other health sector agencies support the following advice.

Stop drinking alcohol if you could be pregnant, are pregnant or are trying to get pregnant. There is no known safe level of alcohol consumption during pregnancy.

A safe level of alcohol consumption during pregnancy has not been established

Regular heavy alcohol use by mothers – defined as more than ten drinks per week – increases the risk of harm^{10 17-19}.

However the association between low to moderate alcohol use (average one drink per day or less) or occasional binge episodes (five or more drinks on a single occasion) and harm is less clear. Some studies have shown increased risk of miscarriage and low birth weight at low levels of alcohol exposure²⁰⁻²², and increased behavioural problems and attention deficits in children exposed to binge drinking^{4 23}. Other studies,

though, have found no deficits in children exposed to low level alcohol exposure or occasional binge drinking in pregnancy^{17 24-28}.

The relationship between maternal drinking and poor child outcomes is complicated by other factors, such as diet, genetics, maternal stress, tobacco smoking, use of marijuana and other substances, as well as the postnatal environment^{29 30}. So not all children exposed to alcohol during pregnancy will be affected or affected to the same degree.

Therefore, given the above, it is not possible to establish a threshold below which alcohol consumption is safe and will not interfere with fetal development³¹.

There is no known safe time to drink alcohol during pregnancy

Because the brain and central nervous system develop throughout pregnancy, there is no period of time when alcohol consumption is safe.

There are, however, time periods where the risk is greater than other times. The first eight weeks of pregnancy is the most vulnerable time for the fetus and a critical time in the development of organs that are especially vulnerable to alcohol exposure, including eyes, heart, ears, and the central nervous system^{2 7 32}.

Unplanned pregnancies pose a particular risk, as alcohol exposure may occur before the woman knows she is pregnant.

What is the role of health professionals in reducing alcohol consumption in pregnancy?

Health professionals can help reduce drinking in pregnancy by asking pregnant women and women trying to get pregnant about their alcohol use at every visit. However, studies suggest that most health professionals do not ask all pregnant women about their alcohol use or provide advice about the effects of alcohol on fetal development³³⁻³⁵.

Screening for alcohol use can improve health outcomes for both mother and child by:

- supporting mothers to reduce or stop drinking
- identifying at-risk pregnancies earlier which can facilitate diagnosis and support for children prenatally exposed to alcohol³¹
- identifying at-risk drinking patterns in non-pregnant women to help to reduce the risk of alcohol exposure for unplanned pregnancies³⁶. Drinking behaviour prior to pregnancy³⁷ is a strong predictor of alcohol consumption during pregnancy^{16,38}.

Women want and expect to receive advice from health professionals about alcohol. Health professionals are seen as a trusted source, having expert knowledge, and being well-placed to support women in changing their drinking behaviour³⁹. Pregnancy is a key opportunity for health professionals to support women to make long term changes to improve their health during and after pregnancy.

Want more information and resources?

Alcohol Drug Helpline

(0800 787 797, alcoholdrughelp.org.nz or free text 8681).

Alcohol and pregnancy information and resources, including information about the Health Promotion Agency's 'Don't know? Don't drink.' campaign, available from alcoholpregnancy.org.nz

Alcohol and Pregnancy – A practical guide for health professionals (Ministry of Health, 2010), available from <https://www.health.govt.nz/system/files/documents/publications/alcohol-pregnancy-practical-guide-health-professionals.pdf>

Implementing the ABC Alcohol Approach in Primary Care (Royal New Zealand College of General Practitioners and the Health Promotion Agency, 2012), available from <https://www.alcohol.org.nz/sites/default/files/documents/2012%20Implementing%20the%20ABC%20Alcohol%20approach%20in%20Primary%20Care%20CEM.pdf>

Information about FASD and the Fetal Alcohol Network NZ available from fan.org.nz

References

1. Gupta KK, Gupta VK, Shirasaka T. An update on Fetal Alcohol Syndrome—Pathogenesis, risks, and treatment. *Alcoholism: Clinical and Experimental Research* 2016;40(8):1594-602.
2. Guerri C, Pascual M. Chapter 24 - Effects of alcohol on embryo/fetal development. In: Gupta RC, ed. *Reproductive and Developmental Toxicology* (Second Edition). Cambridge: Academic Press 2017:431-45.
3. Ghazi Sherbaf F, Aarabi MH, Hosein Yazdi M, et al. White matter microstructure in fetal alcohol spectrum disorders: A systematic review of diffusion tensor imaging studies. *Human Brain Mapping* 2019;40(3):1017-36.
4. Sayal K, Heron J, Golding J, et al. Binge pattern of alcohol consumption during pregnancy and childhood mental health outcomes: Longitudinal population-based study. *Pediatrics* 2009;123(2):e289-96.
5. Falgreen Eriksen HL, Mortensen EL, Kilburn T, et al. The effects of low to moderate prenatal alcohol exposure in early pregnancy on IQ in 5-year-old children. *BJOG: An International Journal of Obstetrics & Gynaecology* 2012;119(10):1191-200.
6. Zuccolo L, Lewis SJ, Smith GD, et al. Prenatal alcohol exposure and offspring cognition and school performance. A 'Mendelian randomization' natural experiment. *International Journal of Epidemiology* 2013;42(5):1358-70.
7. Alvik A, Aalen OO, Lindemann R. Early fetal binge alcohol exposure predicts high behavioral symptom scores in 5.5-year-old children. *Alcoholism: Clinical and Experimental Research* 2013;37(11):1954-62.
8. Harding K, Flannigan K, McFarlane AA. Policy Action Paper: Towards a standard definition of Fetal Alcohol Spectrum Disorder in Canada. Canada: Canada Fetal Alcohol Spectrum Disorder Research Network, 2019.
9. Ministry of Health. Fetal Alcohol Spectrum Disorder 2018 [updated 10.9.2018]. Available from: <https://www.health.govt.nz/our-work/diseases-and-conditions/fetal-alcohol-spectrum-disorder> accessed 2.7.2019.
10. Patra J, Bakker R, Irving H, et al. Dose-response relationship between alcohol consumption before and during pregnancy and the risks of low birthweight, preterm birth and small for gestational age (SGA) - A systematic review and meta-analyses. *BJOG: An International Journal of Obstetrics & Gynaecology* 2011;118(12):1411-21.
11. Irner TB, Teasdale TW, Nielsen T, et al. Substance use during pregnancy and postnatal outcomes. *Journal of Addictive Diseases* 2012;31(1):19-28.
12. Sundermann AC, Zhao S, Young CL, et al. Alcohol use in pregnancy and miscarriage: A systematic review and meta-analysis. *Alcoholism: Clinical and Experimental Research* 2019;43(8):1606-16.
13. Bailey BA, Sokol RJ. Prenatal alcohol exposure and miscarriage, stillbirth, preterm delivery, and sudden infant death syndrome. *Alcohol Research & Health* 2011;34(1):86-91.
14. Ministry of Health. Alcohol use 2012/13 New Zealand Health Survey. Wellington: Ministry of Health, 2015.
15. Rossen F, Newcombe D, Parag V, et al. Alcohol consumption in New Zealand women before and during pregnancy: Findings from the Growing Up in New Zealand study. *The New Zealand Medical Journal* 2018;131(1479):7642.
16. Cheung J, Timmins J, Wright C. Patterns and dynamics of alcohol consumption during pregnancy in a recent New Zealand cohort of expectant mothers. Wellington: Social Policy Evaluation and Research Unit, 2015.

17. O'Keeffe LM, Kearney PM, Greene RA, et al. Maternal alcohol use during pregnancy and offspring trajectories of height and weight: A prospective cohort study. *Drug and Alcohol Dependence* 2015;153:323-9.
18. O'Leary C, Bower C. Guidelines for pregnancy: What's an acceptable risk, and how is the evidence (finally) shaping up? *Drug and Alcohol Review* 2012;31(2):170-83.
19. Strandberg-Larsen K, Poulsen G, Bech BH, et al. Association of light-to-moderate alcohol drinking in pregnancy with preterm birth and birth weight: Elucidating bias by pooling data from nine European cohorts. *European Journal of Epidemiology* 2017;32(9):751-64.
20. Mamluk L, Edwards HB, Savovic J, et al. Low alcohol consumption and pregnancy and childhood outcomes: Time to change guidelines indicating apparently 'safe' levels of alcohol during pregnancy? A systematic review and meta-analyses. *BMJ Open* 2017;7:e015410.
21. Feodor Nilsson S, Andersen PK, Strandberg-Larsen K, et al. Risk factors for miscarriage from a prevention perspective: A nationwide follow-up study. *BJOG: An International Journal of Obstetrics and Gynaecology* 2014;121(11):1375-84.
22. Meyer-Leu Y, Lemola S, Daeppen JB, et al. Association of moderate alcohol use and binge drinking during pregnancy with neonatal health. *Alcoholism: Clinical & Experimental Research* 2011;35(9):1669-77.
23. Niclasen J, Nybo Andersen AM, Teasdale TW, et al. Prenatal exposure to alcohol, and gender differences on child mental health at age seven years. *Journal of Epidemiology and Community Health* 2014;68(3):224-32.
24. Avalos LA, Roberts SCM, Kaskutas LA, et al. Volume and type of alcohol during early pregnancy and the risk of miscarriage. *Substance Use & Misuse* 2014;49(11):1437-45.
25. Lund IO, Moen Eilertsen E, Gjerde LC, et al. Is the association between maternal alcohol consumption in pregnancy and pre school child behavioural and emotional problems causal? Multiple approaches for controlling unmeasured confounding. *Addiction* 2019;114(6):1004-14.
26. McCormack C, Hutchinson D, Burns L, et al. Maternal and partner prenatal alcohol use and infant cognitive development. *Drug and Alcohol Dependence* 2018;185:330-8.
27. Skogerboe A, Kesmodel US, Wimberley T, et al. The effects of low to moderate alcohol consumption and binge drinking in early pregnancy on executive function in 5-year-old children. *BJOG: An International Journal of Obstetrics & Gynaecology* 2012;119(10):1201-10.
28. Underbjerg M, Kesmodel US, Landro NI, et al. The effects of low to moderate alcohol consumption and binge drinking in early pregnancy on selective and sustained attention in 5-year-old children. *BJOG: An International Journal of Obstetrics & Gynaecology* 2012;119(10):1211-21.
29. Murphy DJ, Mullally A, Cleary BJ, et al. Behavioural change in relation to alcohol exposure in early pregnancy and impact on perinatal outcomes - A prospective cohort study. *BMC Pregnancy and Childbirth* 2013;13(no pagination)
30. National Health and Medical Research Council. Australian guidelines to reduce health risks from drinking alcohol. Canberra: National Health and Medical Research Council, 2009.
31. Carson G, Cox LV, Crane J, et al. Alcohol use and pregnancy consensus clinical guidelines. *Journal of Obstetrics and Gynaecology Canada* 2010;32(8, Supplement 3):S1-S2.
32. Ministry of Health. Taking action on Fetal Alcohol Spectrum Disorder (FASD): A discussion document. Wellington: Ministry of Health, 2015.
33. British Medical Association. Alcohol and pregnancy - Preventing and managing fetal alcohol spectrum disorders. London: British Medical Association, 2016.
34. Elek E, Harris SL, Squire CM, et al. Women's knowledge, views, and experiences regarding alcohol use and pregnancy: Opportunities to improve health messages. *American Journal of Health Education* 2013;44(4):177-90.
35. Elliott EJ, Payne J, Haan E, et al. Diagnosis of foetal alcohol syndrome and alcohol use in pregnancy: A survey of paediatricians' knowledge, attitudes and practice. *Journal of Paediatrics and Child Health* 2006;42(11):698-703.
36. Hanson JD, Nelson ME, Jensen JL, et al. Impact of the CHOICES intervention in preventing alcohol-exposed pregnancies in American Indian Women. *Alcoholism: Clinical and Experimental Research* 2017;41(4):828-35.
37. Ministry of Health. Annual update of key results 2017/18: New Zealand Health Survey. Wellington: Ministry of Health, 2019.
38. Skagerström J, Chang G, Nilsen P. Predictors of drinking during pregnancy: A systematic review. *Journal of Women's Health (Larchmt)* 2011;20(6):901-13.
39. Peardon E, Payne J, Parsons, Henley N et al. Alcohol and pregnancy: Women's knowledge, attitudes and practice. *Journal of Paediatrics and Child Health* 2007;43(7-8):A12.